

Socially transmitted diffusion of a novel behaviour from subordinate chimpanzees

Stuart K Watson¹, Lisa A Reamer², Mary Catherine Marenò², Gillian Vale^{1,2}, Rachel A Harrison¹, Susan P Lambeth², Steven J Schapiro^{2,3}, Andrew Whiten*¹

Affiliations

¹Centre for Social Learning and Cognitive Evolution, and Scottish Primate Research Group, School of Psychology and Neuroscience, University of St Andrews, UK

²National Center for Chimpanzee Care, Department of Veterinary Sciences, Michale E. Keeling Center for Comparative Medicine and Research, UT MD Anderson Cancer Center, Bastrop, TX USA

³Department of Experimental Medicine, University of Copenhagen, Denmark

*Corresponding Author

Short title: Subordinate chimpanzees seed traditions

Research highlights:

- Behaviours seeded by subordinate (but not dominant) chimpanzees were copied by observers.
- This finding contrasts with prior work suggesting social learning bias towards dominant chimpanzees.

Contact details for corresponding author

Postal address: Professor Andrew Whiten, School of Psychology & Neuroscience, St Mary's Quad, South Street, St Andrews, KY16 9JP, United Kingdom
Email Address: aw2@st-andrews.ac.uk
Phone: +44 (0)1334 462073

ABSTRACT

Chimpanzees (*Pan troglodytes*) demonstrate much cultural diversity in the wild, yet a majority of novel behaviours do not become group-wide traditions. Since many such novel behaviours are introduced by low-ranking individuals, a bias toward copying dominant individuals ('rank-bias') has been proposed as an explanation for their limited diffusion. Previous experimental work showed that chimpanzees (*Pan troglodytes*) preferentially copy dominant over low-rank models. We investigated whether low ranking individuals may nevertheless successfully seed a beneficial behaviour as a tradition if there are no 'competing' models. In each of four captive groups, either a single high-rank (HR, $n=2$) or a low-rank (LR, $n=2$) chimpanzee model was trained on one method of opening a two-action puzzle-box, before demonstrating the trained method in a group context. This was followed by eight hours of group-wide, open-access to the puzzle-box. Successful manipulations and observers of each manipulation were recorded. Barnard's exact tests showed that individuals in the LR groups used the seeded method as their first-choice option at significantly above chance levels, whereas those in the HR groups did not. Furthermore, individuals in the LR condition used the seeded method on their first attempt significantly more often than those in the HR condition. A network-based diffusion analysis revealed that the best supported statistical models were those in which social transmission occurred *only* in groups with subordinate models. Finally, we report an innovation by a subordinate individual that built cumulatively on existing methods of opening the puzzle-box and was subsequently copied by a dominant observer. These findings illustrate that chimpanzees are motivated to copy rewarding novel behaviours that are demonstrated by subordinate individuals and that, in some cases, social transmission may be constrained by high-rank demonstrators.

Keywords: Social learning, rank, dominance, chimpanzee, culture

INTRODUCTION

It is now generally accepted that social learning is widespread in the animal kingdom and that socially transmitted traditions ('cultures') are found in a wide range of vertebrates [Whiten, 2005; Laland & Janik, 2006; Laland & Galef, 2009]. However, the processes by which a novel behaviour propagates to become a group-wide tradition remain unclear [Rendell et al., 2011]. Indiscriminately copying the behaviours of conspecifics is often not an optimal strategy, as the learner runs the risk of copying costly behaviours or wasting energy on those that are not productive [Kendal, Coolen, van Bergen & Laland, 2005; Rendell et al., 2010]. Accordingly, a number of adaptive 'biases' in social learning have been proposed as possible influences on whether individuals choose to utilise social information and who they get it from, for example 'when uncertain, copy the majority' [Henrich & McElreath, 2003; Laland, 2004; Claidière & Whiten, 2012; van Leeuwen & Haun, 2014]. Due to their cultural diversity [Whiten et al. 1999] and propensity for social learning, chimpanzees have been a favoured model species for studying these social learning biases. Chimpanzees also present an interesting paradox in that although innovations are not an uncommon occurrence, at one field site where researchers made an attempt to quantify their fate it was found that a majority of innovations failed to become group-wide traditions [Nishida, Matsusaka & McGrew, 2009]. The factors that determine whether a novel behaviour diffuses throughout a group or remains limited to one or a minority of individuals are largely unknown. The direct pay-off of a behaviour does not seem sufficient to explain this, given reported instances of the spread of 'arbitrary' traditions with no apparent functional benefits. A striking example of this is described by van Leeuwen, Cronin & Haun [2014], who report a single chimpanzee placing a piece of grass in its ear to no discernible benefit - a 'fashion' which was soon adopted by the rest of the group. Conversely, Hopper et al. [2011] found in a token-exchange task that most chimpanzees chose the same tokens as those selected by a trained model, even

when the alternative token choice resulted in a more preferred food reward, presenting an interesting example of copying a behaviour which is visibly less beneficial than alternatives.

Many novel behaviours enter both wild and captive chimpanzee communities through the lower end of the dominance hierarchy – whether this be from subordinate innovators [Reader & Laland, 2001] or migrant females importing their native behavioural repertoire to their host group [Nakamura & Uehara, 2004; O'Malley, Wallauer, Murray & Goodall, 2012]. A bias toward copying dominant over subordinate individuals has been shown and proposed to explain the relative rarity of these novel behaviours becoming traditions [Kendal et al., 2015]. One might suppose that this would occur for strategic reasons (dominant individuals are successful, so copying them might be an adaptive option), due to normative effects (copying the dominant individual facilitates social cohesion) or simply as a result of an attentional bias towards these individuals (e.g. dominant individuals are central in the social network). In capuchin monkeys it has been found that subordinate individuals tend not to demonstrate acquired token-exchange behaviours in a group context [Addessi et al. 2011] or in the presence of a dominant individual [Lonsdorf et al., 2016], which means there is an inherent rank-bias in the source of social information available to observers. Although it has also been found that capuchins preferentially observe older, more dominant and more proficient nut-crackers in the wild, suggesting a more active learning bias [Coelho et al., 2015]. One or all of these may play a part in restricting the flow of social information from subordinate individuals and cause a group-wide convergence on the behaviour of dominant individuals. To date, two studies have offered evidence for a rank-bias in chimpanzees. Kendal et al. [2015] seeded a method of opening a two-action puzzle box into two groups of chimpanzees using mid-ranking female models (and allowed two other groups to explore the task without trained models), and through complex analysis of attention states during demonstrations found

evidence that individuals preferentially attend to dominant and/or knowledgeable demonstrators. Horner et al. [2010] also concluded that when presented with demonstrations from both a 'high prestige' (high rank and track record as a model) and 'low prestige' (low rank) individual on a token-exchange task, chimpanzees preferentially copied the method demonstrated by the high prestige individual [Horner et al., 2010]. However, there remains the question of whether or not low-ranking individuals, who demonstrate a productive novel behaviour, will be copied if there are no more dominant models available. This question is important for our understanding of how innovations become traditions, and how traditions proliferate across communities.

Accordingly, we compared the diffusion of alternative methods of opening a two-action puzzle-box seeded by either a low- (female) or high-ranking (male) individual in four different groups of chimpanzees. In this context, based on prior work indicating a rank-bias in chimpanzee social learning, we predicted that either (a) social transmission of the seeded method will *only* occur in the groups with high-ranking models (we shall call this the '*hard* rank-bias hypothesis'), or (b) behaviour will be socially transmitted in *both* conditions, but the effect will be stronger in groups with high-ranking models ('*soft* rank-bias hypothesis').

METHODS

Study Site

This study was carried out at the National Center for Chimpanzee Care (NCCC) located at the Michale E. Keeling Center for Comparative Medicine and Research of The University of Texas MD Anderson Cancer Center in Bastrop (UTMDACC), Texas. Data was collected between April and August, 2015. A total of 38 chimpanzees (21 female) participated in the study, aged from 13 to 53 years of age. Most individuals were captive-born, but some (n=5) were wild-born. All individuals have participated in a

wide range of previous behavioural research studies, some of which included puzzle-box tasks, but we have designed our apparatus to require different manipulations to those of earlier studies, as noted below. The participants include both nursery-reared and mother-reared individuals. Following previous studies [Horner et al. 2010, Kendal et al., 2015; Hopper et al., 2015a], the social rank of each individual was determined by surveying the judgments of 5 staff members (behavioural researchers, trainers and management) who had been working with these animals for at least 5 years each. Freeman et al. [2013] found that human assessment of dominance in chimpanzees has good predictive validity for relevant behavioural measures of dominance such as aggression and displacement. Each staff member was asked to rank the individuals in the group linearly from '1' (highest rank) to N (lowest rank) without discussing their rankings with other staff. Agreement between observers was high (>80%), but where disagreements occurred the mode rank for each individual was used. From these rankings we determined the 'alpha' male for each group in the HR condition and chose a subordinate (averaging in the lower third of the hierarchy) female to act as the model for each group in the LR condition. All groups have access to two or more den areas (14m² each) and either an outdoor habitat or dome (dome: 90m², habitat: 400m²) with a range of enrichment devices and activities, and a variety of climbing and swinging structures to promote species-typical behaviours. Testing generally occurred indoors, but access to outdoor enclosures was not restricted. The full demographic and housing information for each participating individual can be found in Table 1. Ethical approval for this study was granted by the School of Psychology & Neuroscience at the University of St Andrews and the IACUC of UTMDACC, adhering to all the legal requirements of US law and the American Society of Primatologists' principles for the ethical treatment of non-human primates. All subjects voluntarily participated in the testing procedures.

ID	CONDITION	SEX	WILD BORN?	DOB	REARING	HOUSING
RAD	LR (1)	M	N	14/01/1990	MOTHER	HABITAT
ANG*	LR (1)	F	Y	01/01/1975	UNKNOWN	HABITAT
CHE	LR (1)	F	N	09/12/1990	NURSERY	HABITAT
KIH	LR (1)	F	N	06/08/1988	NURSERY	HABITAT
MAH	LR (1)	M	N	26/10/1988	MOTHER	HABITAT
NAH	LR (1)	F	N	04/07/1990	NURSERY	HABITAT
AKI	LR (2)	M	N	09/02/1980	NURSERY	DOME
CEC*	LR (2)	F	N	24/02/1991	MOTHER	DOME
HAA	LR (2)	M	N	30/12/1991	MOTHER	DOME
MAR	LR (2)	F	Y	01/01/1966	UNKNOWN	DOME
MART	LR (2)	F	Y	01/01/1965	UNKNOWN	DOME
TAS	LR (2)	F	N	18/11/1992	MOTHER	DOME
ZOE	LR (2)	F	N	13/04/2002	MOTHER	DOME
BRI	HR (1)	F	N	31/08/1995	MOTHER	DOME
CHI	HR (1)	M	N	25/08/1988	MOTHER	DOME
MAG	HR (1)	M	N	24/07/1992	MOTHER	DOME
MAN	HR (1)	F	N	08/09/1984	MOTHER	DOME
NIC*	HR (1)	M	N	07/04/1988	MOTHER	DOME
BER	HR (2)	F	N	18/02/1978	NURSERY	DOME
JUD*	HR (2)	M	N	26/08/1990	NURSERY	DOME
KOB	HR (2)	M	Y	01/01/1972	UNKNOWN	DOME
QUI	HR (2)	F	Y	01/01/1971	UNKNOWN	DOME
TUL	HR (2)	F	N	01/05/1980	MOTHER	DOME
TOD	CONTROL	F	Y	01/01/1971	UNKNOWN	DOME
SAB	CONTROL	F	Y	01/01/1968	UNKNOWN	DOME
PEP	CONTROL	F	Y	01/01/1967	UNKNOWN	DOME
ALP	CONTROL	F	N	08/11/1984	MOTHER	DOME
BET	CONTROL	F	N	23/06/1994	MOTHER	DOME
BIL	CONTROL	M	N	16/06/1993	MOTHER	HABITAT
BO	CONTROL	M	N	16/05/1993	MOTHER	HABITAT
JOE	CONTROL	M	Y	01/01/1972	UNKNOWN	DOME
MAY	CONTROL	F	Y	01/01/1965	UNKNOWN	DOME
MOO	CONTROL	M	Y	01/01/1971	UNKNOWN	DOME
GRE	CONTROL	M	Y	01/01/1970	UNKNOWN	DOME
AJA	CONTROL	M	N	01/01/1978	UNKNOWN	DOME
LUL	CONTROL	F	N	16/01/1982	MOTHER	DOME
TAB	CONTROL	M	N	25/08/1991	MOTHER	DOME
KUD	CONTROL	M	N	07/12/1982	MOTHER	DOME

162 Table 1. Demographic information for all participating individuals. Condition: HR = High

163 rank model, LR = Low-rank model. Asterisk next to name indicates individual was the

164 trained model for their group.

165

166

167

Apparatus

This study employed a two-action, sliding-door puzzle-box (the 'Vert', see Figure 1), a vertical variation we designed to require different actions to those common to earlier social learning studies [Aplin et al., 2015; Hopper, Lambeth, Schapiro & Whiten, 2008; Kendal et al. 2015].




Figure 1. The 'vertical artificial vegetable' (the 'Vert') could be opened to reveal a food reward either by sliding the door entirely upwards (B) or entirely downwards (C). The resting position on presentation is shown in (A). The side-profile is shown in (D). Upon a completed opening, the door locked so as to restrict access to the alternative reward. The anchor platform was attached to a trolley with vice clamps.

Sessions were recorded using a Panasonic HC-X920 video camera. All videos were coded using BORIS, version 2.05 ([www.http://penelope.unito.it/boris](http://penelope.unito.it/boris)). All analyses were carried out using R Statistical Package Version 3.2.3 [2015] with R Studio Version 0.99.491 [2015].

Procedure

For Condition LR (low-rank), in two groups (n=6, 7), a low-ranking female individual was voluntarily separated and trained to open the door by either sliding the door up or down. Likewise in the HR (high-rank) condition, the dominant male of each of two groups (n = 5, 5) was trained on a method of opening the Vert. Females were used for Condition LR and males for Condition HR to maximise the rank disparity between these individuals. Since males are almost always of higher social rank than females in chimpanzee communities, in some groups it would not have been possible to select a high-ranking female to act as a model. However, Kendal et al. [2015] found no bias in whether male or female chimpanzees were preferentially attended to during their experiments, so we would not anticipate sex acting as a confound here. Nevertheless, below we include an analysis of audience sizes during demonstrations of the present study in order to explore whether males and females may differently tolerate observers. LR and HR conditions differed only in the choice of model.

Training began by presenting the baited Vert to the test subject with one of the slide-directions locked so it could not be used. Once a reward had been retrieved successfully 10 times in a row, the alternative method was unlocked and baited for all further trials. Models were considered to be 'trained' once they completed a total of 30 sequential uses of the trained method without deviation.

After being trained, the model was reintroduced to the group and given access to the Vert in a group context. Two 20-minute demonstration sessions were carried out

on subsequent days, during which only the model had access to the box. The Vert was gently pulled out of reach if another individual displaced the model. This was to ensure a roughly equal number of demonstrations between dominant and subordinate models and make the methods comparable with previous work on rank-bias and social learning [Horner et al. 2010; Kendal et al. 2015; Hopper et al. 2015b].

The demonstration period was followed by 8 hours of open-diffusion in which unrestricted access to the Vert was provided. Open-diffusion occurred across multiple sessions, typically of 60 minutes but varying between 45 minutes (due to unforeseen interruptions) and 120 minutes in length (group HR2 had an unavoidably condensed test period, resulting in longer sessions to make up time).

Once any individual in the demonstration or open-diffusion phase had retrieved a reward, the Vert was withdrawn one metre, the door was reset and the reward chamber re-baited. When re-setting the door, the Vert was covered with a cloth to avoid possible directional cues from the experimenter.

To determine whether an inherent directional bias may have influenced which method individuals from experimental groups chose to use, 15 individuals were selected from non-experimental groups to participate in an asocial control condition. Individuals were selected based on advice from care staff about their willingness to voluntarily separate from the group and engage with research procedures. Individuals separated voluntarily from their group and were then presented with the Vert for a period of 20 minutes each. Both reward chambers were baited and both methods of opening the door were unlocked. If an individual completed a successful manipulation of the Vert, the Vert was reset and baited as described above.

233 *Statistical analyses*

234 We used binomial tests to determine whether the number of individuals in the
235 control condition to use each method on their first trial differed significantly from chance
236 (50%), which would indicate an inherent directional bias that would have acted as a
237 confound. We then used Barnard's exact test, an alternative to Fisher's exact test with
238 greater power for small sample sizes [Mehta & Senchaudhuri, 2003], to test whether
239 individuals from high or low rank conditions were significantly more likely to use the
240 seeded method on their first successful trial. Binomial tests were subsequently used to
241 determine whether the proportion of individuals in each condition who used the seeded
242 method on their first successful trial differed significantly from chance (50%). Finally,
243 we applied the same tests to a more conservative, truncated form of the experimental
244 data set. In order to mitigate the possibility that individuals had learned from individuals
245 not of direct interest to the research question, for example a dominant female who had
246 asocially learned the same method as the subordinate model, we only analysed data
247 (for this analysis only) from individuals in both conditions who had *only* observed their
248 group's model demonstrating. This resulted in 11 individuals being excluded from this
249 model, leaving n=8. We also carried out Bayesian equivalents of the analyses
250 described above, which can be found in the Supplemental Material by an interested
251 reader and which were consistent with the findings reported below.

253 *Network-based diffusion analysis*

254 Network-based diffusion analysis (NBDA) is a powerful method of determining
255 whether an observed pattern of acquisition of behaviours is consistent with the
256 predictions of a group's social network [Franz & Nunn, 2009; Hoppitt, Boogert &
257 Laland., 2010; Allen, Weinrich, Hoppitt & Rendell, 2013; Hobaiter, Poisot, Zuberbuhler,
258 Hoppitt & Gruber, 2014]. In this case, the social network was created using the number

of times Individual A observed Individual B using the seeded method before Individual A first demonstrated this method. Because we were able to record the exact times at which an individual first used the method, we used the Time of Acquisition Diffusion Analysis (TADA) variant of NBDA [Hoppitt et al., 2010]. Times entered into the model were the number of seconds which the group had been exposed to the Vert before a given individual first opened it using the seeded method.

We used an information theoretic approach [Burnham & Anderson, 2002], using Akaike's information criterion corrected for sample size (AICc) from which total Akaike weights ($\sum w_i$) for each model were calculated. Total Akaike Weights were then used to create model averaged estimates for the factor by which individuals' learning rates are increased per observation of the seeded method. Models were constructed based on the predictions outlined by the rank-bias hypothesis and the necessary conditions for refutation (above).

This analysis was carried out using the NBDA R Script Version 1.2.11 (available at <http://lalandlab.st-andrews.ac.uk/freeware/>).

Generalised linear mixed effects models

We used two sets of generalised linear mixed effects models (GLMMs) to determine whether the sex of a demonstrator was a useful predictor in determining how many individuals were likely to be in proximity (<3m) on any given trial. The first set of models considered audience size as an absolute value, whereas the second considered it as a proportion of group size. In all models, 'individual' was fit with random intercepts and random slopes to account for multiple measurements from each individual. We took an information theoretic approach to inference, using akaike's information criterion corrected for small sample sizes (AICc) to estimate model fit. From this we calculated total akaike weights ($\sum w_i$) and use these to compute model-averaged

estimates of parameter coefficients, allowing us to estimate the effect of a parameter while taking into account model uncertainty. Due to the use of model-averaging, rather than use p-values to determine whether a parameter had an important effect on the output variable, this was established according to whether its 95% confidence intervals overlapped with 0.


Video Coding

The method used by any individual who successfully opened the box was coded, as well as the identities of any individual within 3 meters. Any individuals within 3m whose heads were oriented towards the Vert and did not have their view obstructed was recorded as having observed the opening. Videos were coded by SKW. Inter-observer reliability was carried out with RAH on the method ('up' or 'down') used and who was observing each demonstration in 30 clips of individuals opening the Vert, with 100% agreement.

RESULTS

Although the raw data from the control condition (Figure 2) are somewhat suggestive of a greater tendency for pushing down than lifting up the door, the number of individuals who chose either method did not differ significantly from chance (Up: $n = 3$, $P = 0.343$, 95% CI = 0.07-0.65; Down: $n = 7$, $P = 0.343$, 95% CI: 0.35-0.93). Nevertheless, the direction of the seeded method was counterbalanced across groups in the experimental condition. Furthermore, five out of 15 control individuals failed to open the box at all, from which we may infer that the two methods of opening the door were not so salient that every chimpanzee was easily capable of opening it without the use of social information.

310




311

312 Figure 2. Method used on first opening of the Vert by individuals in the control
313 condition, and number of individuals who failed to open the box.

314

315 A Barnard's exact test found that individuals in the low-rank condition used the
316 seeded method on their first successful trial significantly more often than individuals in
317 the high-rank condition ($X^2 = 2.09$, $N=19$, $P=0.048$, see Figure 3). Exact binomial tests
318 found that individuals in the low-rank condition used the seeded method significantly
319 more often than chance ($n = 11$, $P=0.032$, 95% CI=0.53-1.0) but high-rank condition
320 did not ($n=8$, $P=0.855$, 95% CI=0.111 – 1.0).



321

322 Figure 3. Column 1 - Methods used on first opening of the box for each condition.

323 Columns 2 and 3 - Methods used on first opening of the box in each group.

324 Directionality of trained method indicated for each group in brackets.

325

326 Using a truncated data set (Figure 4: procedure and rationale for exclusion
 327 detailed above), there remained a significant difference between low and high-rank
 328 conditions in the number of individuals who used the seeded method on their first trial
 329 (Barnard's exact test: $X^2=2.19$, $n=8$, $P=0.047$). However, it is worth noting that two of
 330 the four individuals in HR condition (see HR2 in Table 2) who first used the non-seeded
 331 method later switched to consistently use the seeded method. No other individuals in
 332 any group persistently switched to a method other than their first-learned, with the
 333 exception of the innovation described in detail below. Both individuals who did not
 334 solve the task were males. Neither individual tried any other method of interacting with
 335 the door (e.g. hitting, pushing, pulling, etc), indicating that they lacked the motivation to
 336 engage with the task.




Figure 4. Counts for methods used on first opening of the box in the original ‘full’ data set, side-by-side with ‘truncated’ data set.

ID	Group	First method as seeded?	Total trials	Total trials as seeded
JUD	HR1	Y	730	720
BER	HR1	N	69	0
TUL	HR1	Y	234	214
QUI	HR1	N	1	0
KOB	HR1	n/a	0	0
NIC	HR2	Y	535	463
CHI	HR2	N	109	108
MAN	HR2	N	54	48
MAG	HR2	Y	106	66
BRI	HR2	Y	9	6
CEC	LR1	Y	170	170
MAY	LR1	Y	185	184
ZOE	LR1	Y	123	121
AKI	LR1	N	171	3
TAS	LR1	Y	166	163
MAR	LR1	Y	34	34
HAA	LR1	Y	138	138
ANG	LR2	Y	146	146
CHE	LR2	Y	115	108
KIH	LR2	Y	326	133
NAH	LR2	Y	188	162
RAD	LR2	Y	13	13
MAH	LR2	n/a	0	0

Table 2. Summary table of each individual’s interactions with the puzzle box.

Network-based Diffusion Analysis

There was most support for models (Table 3) in which there was an effect of social transmission (S) in the LR condition but not HR, with S varying between groups ($\Sigma w_i = 0.75$). Model-averaged estimates for S indicate that each observation increased an average individual's learning rate by 3% in LR1 and 15% in LR2. Model averaged estimates for S indicate that each observation increased an average individual's learning rate in HR1 and HR2 by 0.1% per observation. Models based on the hard rank-bias hypothesis were not well supported ($\Sigma w_i = 0.002$ and $\Sigma w_i = 0.009$). A model allowing for the soft rank-bias hypothesis had some support ($\Sigma w_i = 0.078$), but contrary to the predictions of this hypothesis, the effect of S was estimated as being greater in the LR condition ($S = 0.08$) than HR ($S = 0.00$). Individual-level variables (sex, age and rearing history) were added to the best fitting model, but there was little support for any of them improving the model (Table 4).

Model	AICc	Delta AICc	Total weighted AICc (Σw_i)
*S only in HR, varies between HR groups	334.13	11.91	0.002
S in all groups	330.96	8.74	0.009
*S only in HR, constant between HR groups	330.96	8.74	0.009
S varies between all groups	328.6	6.38	0.027
No S in any group	328.27	6.05	0.036
**S varies between LR and HR	326.75	4.53	0.078
S only in LR, constant between LR groups	326.5	4.28	0.088
S only in LR, varies between LR groups	322.22	0	0.75

* 'Hard' rank bias hypothesis

** 'Soft' rank-bias hypothesis candidate

Table 3. AICc, delta AICc and Total Akaike Weights (Σw_i) for each model. 'S' = social transmission.

Asocial				
variable	df	AICc	Delta AICc	Total weighted AICc
Sex	4	325.73	3.51	0.07
Rearing	4	323.13	0.91	0.25
Age	4	323.08	0.86	0.26
None	3	322.22	0	0.41

Table 4. AICc, delta AICc and Total Akaike Weights (Σw_i) for the best fitting model from Table 3 with additional individual-level variables.

GLMMs

A model-averaged estimate (Table 5) of the coefficient for the effect of demonstrator sex on audience size when counting absolute number of individuals within 3m was 0.14 (95% CI: -0.23, 0.51), and when considering audience size as a proportion of total group size was 0.02 (95% CI: -0.05, 0.08). We may infer that Sex did not have an important effect as the 95% confidence intervals did not overlap with zero. Furthermore, as seen in Table 5, adding Sex to the models resulted in a considerably higher AICc and therefore poorer fit.

GLMM Set 1: Audience = Number of individuals < 3m from demonstrator

Model	K	AICc	Delta AICc	Total AICc weight
Audience ~ 1 ID	3	9178.19	0	0.94
Audience ~ Sex + Sex ID	6	9183.63	5.44	0.06

GLMM Set 2: Audience = Proportion of group < 3m from demonstrator

Model	K	AICc	Delta AICc	Total AICc weight
Audience ~ 1 ID	3	-2114.53	0	0.94
Audience ~ Sex + Sex ID	6	-2108.95	5.58	0.06

Table 5. Model comparison summary statistics for two sets of GLMMs. Sex = Sex of demonstrator. 1|ID = Random intercepts for individual. Sex|ID = random slopes and intercepts for Individual. K = number of effective parameters.

An Innovation

Finally, we report an innovation which occurred in one of the high-rank condition groups. After 7 hours of open-diffusion, a subordinate individual (TUL) discovered a narrow window of motion in which the door can be opened using 'Up', so that a reward can be retrieved, but the locking mechanism is not activated. This allowed her to then also use 'Down' to move the door a second time and obtain a second reward. TUL had not used 'Down' prior to this discovery, but had observed two other females in her group using it on multiple occasions. This suggests TUL combined her first-learned method with previously acquired social information about that used by others to generate a more productive method, although asocial learning cannot be ruled out. Despite the innovator being of low rank, after 11 observations of this improved method the dominant male (JUD) of the group, who to this point had exclusively used the 'Up' method, also began to use the combined form. A similar pattern was observed in a second group. Again, the first individual was a subordinate female (CHE) and the method was subsequently used by two higher ranking females (KIH, NAH). Due to the limited data available, it is not possible to carry out any formal analyses of these events, but we present them as 'naturally' occurring examples of subordinates' innovations achieving limited diffusion through their groups.

DISCUSSION

Rank-bias has been proposed as a way to account for the relatively rare adoption of innovations to produce traditions within chimpanzee communities [Horner et al., 2010; Kendal et al., 2015]. Based on this 'rank-bias hypothesis', we predicted that novel behaviours seeded by subordinates either fail to spread, or motivate a considerably lesser degree of social learning than novel behaviours seeded by dominant individuals. In our study, not only were the group-mates of low-ranking

models more likely to use the seeded rather than non-seeded method on their first opening of the box, but they were also substantially more likely to do so than individuals in groups with high-rank models. Furthermore, a NBDA showed greatest support for models in which social transmission of the seeded method was present only in the low-rank condition. Finally, we reported innovations developed by two subordinate chimpanzees in separate groups which built on pre-existing methods and were subsequently used by more dominant individuals, likely as a result of social learning. While one must be cautious in interpreting isolated events, these instances are striking in their pertinence to our research question and in how they contrast with the predictions of the rank-bias hypothesis.

We conclude these findings strongly suggest that the rank-bias identified by previous studies [Kendal et al. 2015; or 'prestige-bias' in Horner et al., 2010], which occurred when observers had a choice between models of various ranks, does not prohibit the successful emergence of group-wide behaviour patterns from subordinate models or innovators when no competing model is present. As well as a rank-bias, Kendal et al. [2015] identified a bias towards copying 'knowledgeable' individuals, which our results suggest to be the case even when demonstrators are of low social rank. This may make adaptive sense, since if one observes an individual doing something that is rewarding, it is counterintuitive to ignore this information simply on the basis of the demonstrator's low social status. However, this does not preclude the indirect importance of rank in more natural settings. For example, if recent immigrants tend to be spatially peripheral to the group, this would reduce the number of individuals in close enough proximity to observe (and copy) any novel behaviours being used, functionally resulting in a rank bias. By contrast, individuals in the present study could only carry out the behaviour when performing it in a central, commonly used space where the researcher and experiment were set up, making them readily visible to their group. Furthermore, while we did identify comprehensive diffusion of methods seeded

subordinate models, it is important to note the difference in group size between the relatively small groups studied here (between 6 and 8 individuals) and wild chimpanzee communities which can have anything from 20 to 150 members [Goodall 1986; Nishida 1990; Boesch & Boesch-Achermann 2000]. Communities of larger scale, as well as the presence of fission-fusion social dynamics, may present additional obstacles for behavioural diffusion.

Being raised in captivity and participating in behavioural research for so many years [e.g. Brosnan et al., 2007; Hopper, Lambeth, Schapiro & Whiten, 2008; Dean, Kendal, Schapiro, Thierry & Laland, 2012; Kendal et al., 2015] may also have shaped the study population to be more ready social learners [Carpenter & Tomasello, 1995], further mediating the effects of rank-bias. The influence of such developmental, cultural, environmental and individual differences on social learning are difficult to examine in such long-lived species, but are likely to be critical in our understanding of cultural transmission [Mesoudi, Chang, Dall & Thornton, 2016]. Nevertheless, this would not explain why there was a greater effect of social transmission in the low-rank condition than in our high-rank condition.

These results contrast with prior studies [Horner et al. 2010; Kendal et al. 2015] in that the effect of social transmission was found to be stronger in our low-rank condition, and a greater proportion of individuals in the LR condition used the seeded method on their first trial than those in HR. One methodological difference between the current study and previous work that might explain this discrepancy is that our high-ranking models were dominant males rather than dominant females. This was an intentional design choice, as males are almost always dominant relative to females, and it was desirable to maximise the rank disparity between model types. However, this may have introduced additional confounds. While males were successfully used as models in Price, Lambeth, Schapiro & Whiten [2009], the study used video demonstrations and observers were not always from the same group as the model, and

were therefore unaware of their rank. Wrangham et al. [2016] found that in a community of chimpanzees where multiple grooming techniques were in use, individuals tended to converge on the method primarily used by their matriline, potentially hinting at a sex bias in chimpanzee social learning. However, the only systematically documented example of an incipient tradition diffusing through a wild chimpanzee community originated in a male chimpanzee [Hobaiter et al. 2014], indicating that males can also make effective models. Furthermore, in a series of GLMM's we examined whether the number of individuals in proximity or attending to an individual's demonstrations could be predicted by that demonstrator's sex, and this was not found to be the case (Table 5). From this we may infer that our use of differently sexed models did not introduce an important confound with respect to social tolerance that would explain the contrast between effects of high versus low rank models in our study. In any case, the key finding in our results is not so much the contrast between effects of high versus low ranked models, but that the low ranked female provided an adequate model whose preferred behavioural option was copied by others.

There is already good evidence for an attentional bias toward dominant individuals [Kendal et al. 2015], but it is unclear to what extent this may be vigilance rather than active social learning. Spatial tolerance between demonstrators and observers is also likely to be crucial in facilitating social learning [van Schaik, Fragaszy & Perry, 2003], which may be confounded when highly dominant demonstrators monopolise a resource. The difficulty associated with faithfully copying a socially intolerant individual may explain why two observers in the HR condition first discovered the non-seeded method and then switched to consistently use the seeded method for the remainder of testing. Based on previous work [Hrubesch, Preuschoft & van Schaik, 2009] we would expect such individuals to fixate on their first-learned method, since the alternative did not provide a greater payoff [van Leeuwen et al., 2013]. It may be that,

in this case, the first-used method was an ‘accidental’ discovery on the route to learning the seeded method.

As previously discussed, capuchin monkeys inhibit demonstration of known behaviours while in the presence of dominant males [Lonsdorf et al. 2016]. If the same is true of chimpanzees, then non-dominant individuals having to wait for an appropriate social context to interact with the task may have introduced additional demands on memory that would interfere with accurate copying models in the HR condition. In our experiment, the fact that we removed the Vert when models were displaced in the demonstration phase meant that the resource could not be immediately monopolised. The reason for this was to remain methodologically consistent with prior work on rank-bias [Horner et al. 2010; Kendal et al. 2015], as well as to directly examine the motivation of observers to learn from subordinate models rather than the effects of resource-monopolisation on the diffusion of novel behaviours. Competition over resources remains an unexamined and potentially important influence on the diffusion of chimpanzee traditions.

While this study has shown that chimpanzees are motivated to learn novel methods of accessing a resource from subordinate individuals, it is possible this is not true of forms of imitative behaviour that are thought to be normatively motivated and therefore, perhaps particularly directed toward important social partners. Examples of this include the fashion of putting grass in one’s ear, invented by a high-ranking female, described by van Leeuwen et al. [2014] or vocal convergence resulting from close social affiliation [Fedurek et al. 2013; Watson et al. 2015]. Further examination of context-specific qualities, such as behavioural-domain, extrinsic motivators (e.g food or social benefits), ease of monopolisation and how these inhibit or promote particular learning biases, may be a fruitful area of research.

ACKNOWLEDGEMENTS

SKW, GV, RAH and AW are grateful for the support of the John Templeton Foundation, grant ID40128, 'Exploring the evolutionary foundations of cultural complexity, creativity and trust' to AW and Kevin Laland, which funded this project. Thanks to Will Hoppitt and Charlotte Brand for statistical discussion. Thanks also to the anonymous reviewers whose suggestions greatly improved this manuscript. The National Center for Chimpanzee Care is supported by NIH Cooperative Agreement U42 OD-011197.

REFERENCES

- Allen J, Weinrich M, Hoppitt W, Rendell L. 2013. Network-based diffusion analysis reveals cultural transmission of lobtail feeding in humpback whales. *Science* 340:485-488.
- Addessi E, Mancini A, Crescimbene L, Visalberghi E. 2011. How social context, token value, and time course affect token exchange in capuchin monkeys (*Cebus apella*). *International Journal of Primatology*. 32:83-98.
- Aplin LM, Farine DR, Morand-Ferron J, Cockburn A, Thornton A, Sheldon BC. 2015. Experimentally induced innovations lead to persistent culture via conformity in wild birds. *Nature* 518:538-541.
- Boesch C., Boesch-Achermann H. 2000. *The chimpanzees of the Tai Forest: Behavioural ecology and evolution*. Oxford University Press, USA.
- Brosnan SF, Jones OD, Lambeth SP, Mareno MC, Richardson AS, Schapiro SJ. 2007. Endowment effects in chimpanzees. *Current Biology* 17:1704-1707.
- Burnham KP, Anderson DR. 2004. Multimodel inference understanding AIC and BIC in model selection. *Sociological methods & research* 33:261-304.

532 Carpenter M, Tomasello, M. 1995. Joint attention and imitative learning in children,
533 chimpanzees, and enculturated chimpanzees. *Social Development* 4:217-237.

534 Claidière N, Whiten A. 2012. Integrating the study of conformity and culture in humans
535 and nonhuman animals. *Psychological bulletin* 138:126.

536 Coelho CG, Falótico T, Izar P, Mannu M, Resende BD, Siqueira JO, Ottoni EB. 2015.
537 Social learning strategies for nut-cracking by tufted capuchin monkeys (*Sapajus*
538 spp.). *Animal cognition*, 18:911-9.

539 Dean LG, Kendal RL, Schapiro SJ, Thierry B, Laland KN. 2012. Identification of the
540 social and cognitive processes underlying human cumulative
541 culture. *Science* 335:1114-1118.

542 Franz M, Nunn CL. 2009. Network-based diffusion analysis: a new method for
543 detecting social learning. *Proceedings of the Royal Society of London B:*
544 *Biological Sciences* 276:1829-1836.

545 Fedurek P, Machanda ZP, Schel AM, Slocombe KE. 2013. Pant hoot chorusing and
546 social bonds in male chimpanzees. *Animal Behaviour* 86:189-196.

547 Friard O, Gamba M. 2016. BORIS: a free, versatile open-source event-logging software
548 for video/audio coding and live observations. *Methods in ecology and Evolution*.
549 DOI: 10.1111/2041-210X.12584

550 Goodall J. 1986. *The chimpanzees of Gombe: Patterns of behavior*. Cambridge, MA:
551 Belknap Press of Harvard University Press

552 Henrich J, McElreath R. 2003. The evolution of cultural evolution. *Evolutionary*
553 *Anthropology: Issues, News, and Reviews* 12:123-135.

554 Hobaiter C, Poisot T, Zuberbühler K, Hoppitt W, Gruber T. 2014. Social network
555 analysis shows direct evidence for social transmission of tool use in wild
556 chimpanzees. *PLoS Biol* 12:e1001960.

557 Hopper LM, Schapiro SJ, Lambeth SP, Brosnan SF. 2011. Chimpanzees' socially
558 maintained food preferences indicate both conservatism and conformity. *Animal*
559 *Behaviour* 81:1195-1202.

560 Hopper LM, Lambeth SP, Schapiro SJ, Whiten A. 2008. Observational learning in
561 chimpanzees and children studied through 'ghost' conditions. *Proceedings of the*
562 *Royal Society of London B: Biological Sciences* 275:835-840.

563 Hopper LM, Kurtycz LM, Ross SR, Bonnie KE. 2015a. Captive chimpanzee foraging in
564 a social setting: a test of problem solving, flexibility, and spatial discounting.
565 *PeerJ*, 17;3:e833.

566 Hopper LM, Lambeth SP, Schapiro SJ, Whiten A. 2015b. The importance of witnessed
567 agency in chimpanzee social learning of tool use. *Behavioural processes*
568 112:120-9.

569 Hoppitt W, Boogert NJ, Laland KN. 2010. Detecting social transmission in
570 networks. *Journal of Theoretical Biology* 263:544-555.

571 Horner V, Proctor D, Bonnie KE, Whiten A, de Waal FB. 2010. Prestige affects cultural
572 learning in chimpanzees. *PLoS One* 5:e10625.

573 Kendal RL, Coolen I, van Bergen Y, Laland, KN. 2005. Trade-offs in the adaptive use
574 of social and asocial learning. *Advances in the Study of Behavior*, 35, 333-379.

575 Kendal RL, Hopper LM, Whiten A, Brosnan SF, Lambeth SP, Schapiro SJ, Hoppitt W.
576 2015. Chimpanzees copy dominant and knowledgeable individuals: implications
577 for cultural diversity. *Evolution and Human Behavior* 36:65-72.

578 Laland KN. 2004. Social learning strategies. *Animal Learning & Behavior* 32:4-14.

579 Laland KN, Janik VM. 2006. The animal cultures debate. *Trends in Ecology &*
580 *Evolution* 21:542-547.

581 Laland KN, Galef BG. 2009. *The question of animal culture*. Harvard University Press.

582 Lonsdorf EV, Bonnie KE, Grim M, Krupnick A, Prestipino M, Whyte J. 2016. Seeding
583 an arbitrary convention in capuchin monkeys: the effect of social context.
584 Behaviour. 153:633-54.

585 Mehta CR, Senchaudhuri P. 2003. Conditional versus unconditional exact tests for
586 comparing two binomials. Cytel Software Corporation, 675.

587 Mesoudi A, Chang L, Dall SR, Thornton A. 2016. The Evolution of Individual and
588 Cultural Variation in Social Learning. Trends in Ecology & Evolution 31:215:225

589 Nakamura M, Uehara S. 2004. Proximate Factors of Different Types of Grooming
590 Hand-Clasp in Mahale Chimpanzees: Implications for Chimpanzee Social
591 Customs 1. Current Anthropology 45:108-114.

592 Nishida T. 1990. *Chimpanzees of the Mahale Mountains*. University of Tokyo Press.

593 Nishida T, Matsusaka T, McGrew WC. 2009. Emergence, propagation or
594 disappearance of novel behavioral patterns in the habituated chimpanzees of
595 Mahale: a review. Primates 50:23-36.

596 O'Malley RC, Wallauer W, Murray CM, Goodall J. 2012. The appearance and spread of
597 ant fishing among the Kasekela chimpanzees of Gombe: a possible case of
598 intercommunity cultural transmission. Current anthropology 53:650.

599 RStudio Team. 2015. RStudio: Integrated Development for R. RStudio, Inc., Boston,
600 MA URL <http://www.rstudio.com/>.

601 Price EE, Lambeth SP, Schapiro SJ, Whiten A. 2009. A potent effect of observational
602 learning on chimpanzee tool construction. Proceedings of the Royal Society of
603 London B: Biological Sciences, rspb20090640.

604 Reader SM, Laland KN. 2001. Primate innovation: sex, age and social rank
605 differences. International Journal of Primatology 22:787-805.

606 Rendell L, Boyd R, Cownden D, Enquist M, Eriksson K, Feldman MW, Fogarty L,
607 Ghirlanda S, Lillicrap T, Laland KN. 2010. Why copy others? Insights from the
608 social learning strategies tournament. *Science* 328:208-13.

609 Rendell L, Fogarty L, Hoppitt WJ, Morgan TJ, Webster MM, Laland KN. 2011.
610 Cognitive culture: theoretical and empirical insights into social learning
611 strategies. *Trends in cognitive sciences* 15:68-76.

612 R Core Team. 2015. R: A language and environment for statistical computing. R
613 Foundation for Statistical Computing, Vienna, Austria. URL [http://www.R-](http://www.R-project.org/)
614 [project.org/](http://www.R-project.org/).

615 Van Leeuwen EJ, Cronin KA, Haun DB. 2014. A group-specific arbitrary tradition in
616 chimpanzees (*Pan troglodytes*). *Animal cognition* 17:1421-1425.

617 Van Leeuwen EJ, Haun DB. 2014. Conformity without majority? The case for
618 demarcating social from majority influences. *Animal Behaviour* 96:187-194.

619 Van Schaik CP, Fragaszy D, Perry S. 2003. Local traditions in orangutans and
620 chimpanzees: social learning and social tolerance. *The biology of traditions:*
621 *Models and evidence* 297-328.

622 Watson SK, Townsend SW, Schel AM, Wilke C, Wallace EK, Cheng L, West V,
623 Slocombe, KE. 2015. Vocal learning in the functionally referential food grunts of
624 chimpanzees. *Current Biology* 25:495-499.

625 Whiten A, Goodall J, McGrew WC, Nishida T, Reynolds V, Sugiyama Y, Tutin CEG,
626 Wrangham RW, Boesch C. 1999. Cultures in chimpanzees. *Nature* 399:682-685.

627 Whiten A. 2005. The second inheritance system of chimpanzees and humans. *Nature*
628 437:52-55.

SUPPLEMENTAL MATERIAL

Bayesian analysis

For the interested reader, we present Bayesian equivalents to the statistical analyses carried out in the main text of this article. A Bayesian approach to statistical inference has the benefit of being better able to handle small data sets that are not normally distributed [Schoot et al., 2014], as is the case in the present study.

Bayesian binomial regression models using MCMC (3 chains of 5,000 iterations, warmup = 1000 iterations) were used for three analyses. Firstly, we determined whether the control group demonstrated any strong directional bias on their first opening of the Vert, which might confound experimental conditions. Secondly, we estimated the effect of condition (HR or LR) on the likelihood that an individual used the seeded method on their first attempt at opening the Vert. The third model used a more conservative, truncated form of the experimental data set. In order to mitigate the possibility that individuals had learned from individuals not of direct interest to the research question, for example a dominant female who had asocially learned the same method as the subordinate model, we only analysed data (for this analysis only) from individuals in both conditions who had *only* observed their group's model demonstrating. This resulted in 11 individuals being excluded from this model, leaving $n=8$. All statistical models described here specified uniform, non-informative priors. Analyses were run using the R package 'Rethinking' version 1.59 [McElreath, 2014] and MCMC was implemented in Rstan (Gelman, 2014) version 2.10.1. Mixing was assessed visually. For each parameter in each model we report the estimated probability of the described event (e.g. using 'Up' on first opening of the box) and the bounds in which there is a 95% probability that the true value lies (95% credible intervals).

Results

A binomial regression model based on the data from individuals in the control condition (see Figure 2) estimated that the probability of individuals using 'up' on their first attempt was 0.22 [95% Credible Interval (CI): 0.08, 0.48], and of using 'down' was 0.47 [95% CI: 0.25, 0.70]. Although these results are somewhat suggestive of a greater tendency for pushing down than lifting up the door during exploration of the box, the wide, overlapping credibility intervals indicate that the bias is not particularly strong. Nevertheless, the direction of the seeded method was counterbalanced across groups in the experimental condition. Furthermore, five out of 15 control individuals failed to open the box at all, from which we may infer that the two methods of opening the door were not so salient that every chimpanzee was easily capable of opening it without the use of social information.

A second model based on our experimental conditions (Figure 3) found an important effect of Condition (Beta = 1.64, 95% CI: 0.07, 3.45) on the likelihood that an individual will use the seeded method on their first opening of the box. The model estimated that the absolute probability that individuals in the HR condition would use the seeded method on their first opening was 0.36 (95% CI: 0.10, 0.70) and 0.91 (95% CI: 0.86, 0.99) for individuals in the LR condition.

This pattern also held when applying the same analysis to the truncated dataset (Figure 4) the rationale and procedure for which was described in the Methods section, above). A main effect of condition (HR/LR) was found (Beta = 10.34, 95% CI: 2.12, 18.76). The estimated probability of individuals in HR using the seeded method on their first opening of the box was 0.018 [95% CI: 0.01, 0.63] in the HR condition, and 0.79 [95% CI: 0.31, 0.98] in the LR condition. However, it is worth noting that two of the four individuals in HR condition (see HR2 in Table 2) who first used the non-seeded method later switched to consistently use the seeded method. No other

54 individuals in any group persistently switched to a method other than their first-learned,
55 with the exception of the innovation described in the main text.

56

57 **References**

58 Gelman, A. (2014). Rstan: the R interface to Stan.

59 McElreath, R. (2016). rethinking: Statistical Rethinking book package. R package
60 version. 1.59.

61 Schoot R, Kaplan D, Denissen J, Asendorpf JB, Neyer FJ, Aken MA. 2014. A gentle
62 introduction to Bayesian analysis: applications to developmental research. Child
63 development 85:842-860.